Feline Picornavirus. Canine Distemper Virus. Avian food or water, utensils, instruments, cages, kennels through the skin. Harmful if swallowed. Wear googles or Influenza A Virus, and Porcine Respiratory & Reproductive stables, catteries, etc. Remove all animals and feed from face shield, rubber gloves, and protective clothing Syndrome Virus (PRRSV), Add 2 ounces of the product per premises, animal transportation vehicles, crates, etc Remove contaminated clothing and wash before reuse. iallon of water. For heavily soiled areas, a pre-cleaning Remove all litter, droppings and manure from floors, walls Wash thoroughly with soap and water after handling and INERT INGREDIENTS: 94.575% step is required. Apply solution with a cloth, mop, sponge, and surfaces occupied or traversed by animals before eating, drinking, or using tobacco hand pump trigger sprayer or low pressure coarse sprayer. Thoroughly clean all surfaces with soap or detergent and so as to wet all surfaces thoroughly. Allow the surface to rinse with water. Saturate surfaces with a use-solution of 2 STORAGE AND DISPOSAL emain wet for 10 minutes, then remove excess liquid. ounces of Lemon Disinfectant Neutral Cleaner - 64 per DO NOT CONTAMINATE WATER, FOOD, OR FEED BY KILLS HIV-1, HIV-2, HBV, and HCV ON PRE- gallon of water for a period of 10 minutes. Ventilate STORAGE OR DISPOSAL CLEANED ENVIRONMENTAL SURFACES/OBJECTS buildings and other closed spaces. Do not house animals PESTICIDE STORAGE - Store in original container in a PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in or employ equipment until treatment has been absorbed, cool (temperature no lower than 50°F or higher than health care settings (Hospitals, Nursing Homes) or other set or dried 120°F), dry and locked storage area inaccessible to ettings in which there is an expected likelihood of soiling Disinfection of Barber/Salon Tools Directions; children. of inanimate surfaces/objects with blood or body fluids, Immerse pre-cleaned barber/salon tools (combs. PESTICIDE DISPOSAL - Pesticide wastes are acutely and in which the surfaces/objects likely to be soiled with brushes, razors, manicure/pedicure tools, clippers, hazardous, Improper disposal of excess pesticide, spray blood or body fluids can be associated with the potential scissors, trimmer blades) in a use-solution of 2 ounces of mixture, or rinsate is a violation of Federal Law. If these for transmission of Human Immunodeficiency Virus Type 1, this product per gallon of water for at least 10 minutes, wastes cannot be disposed of by use according to label advice. or Type 2 (HIV-1 or HIV-2) (associated with AIDS), Hepatitis Rinse instruments thoroughly and dry before reuse. A instructions, contact your State Pesticide or fresh use-solution must be prepared daily or more often if Environmental Control Agency, or the Hazardous Waste SPECIAL INSTRUCTIONS FOR CLEANING AND the use-solution becomes visibly dirty, NOTE: Plastics representative at the nearest EPA Regional Office for DECONTAMINATION AGAINST HIV-1, HIV-2, HBV, may remain immersed until ready to use. Stainless steel guidance and HCV OF SURFACES/OBJECTS SOILED WITH shears and instruments must be removed from the use- CONTAINER DISPOSAL - Nonrefillable container. **BLOOD/BODY FLUIDS.** solution after 10 minutes, rinsed, dried and kept in a clean, not reuse or refill this container. Clean container promptly PERSONAL PROTECTION: When handling items soiled non-contaminated receptacle. Prolonged immersion may after emptying. After rinsing container, offer empty metal with blood or body fluids, use disposable latex gloves, cause damage to stainless steel or metal instruments. drums or plastic containers for recycling if available or GENERAL CLEANING - Apply Lemon Disinfectant puncture and dispose of in a sanitary landfill or by owns, masks, and eye coverings. CLEANING PROCEDURES: Blood and other body fluids Neutral Cleaner - 64 to soiled area with a mop. cloth, incineration. must be thoroughly cleaned from surfaces and objects sponge, hand pump trigger sprayer or low pressure RINSING INSTRUCTIONS before application of this product coarse sprayer. This product can be diluted at 2 ounces Triple rinse container promptly after emptying. Triple rins CONTACT TIME: Allow surface to remain wet for 10 per gallon or applied at full strength. Thoroughly wet as follows: Empty the remaining contents into application Probable mucosal damage may contraindicate the use of gastric layage equipment or a mix tank and drain for 10 seconds after the soiled surface and scrub as necessary. DISPOSAL OF INFECTIOUS MATERIALS: Blood and DEODORIZATION - To deodorize, apply Lemon flow begins to drip. Fill the container ¼ full with water and other body fluids must be autoclaved and disposed of Disinfectant Neutral Cleaner - 64 as indicated under the recap. Shake for 10 seconds. Pour rinsate into application according to local regulations for infectious waste heading DISINFECTION. equipment or a mix tank or store rinsate for later use or MILDEWSTAT - To control mold and mildew (Aspergillus disposal. Drain for 10 seconds after the flow begins to LEMON DISINFECTANT NEUTRAL CLEANER IS A niger) and the odors they cause on pre-cleaned, hard, drip Repeat this procedure two more times. HOSPITAL USE BACTERICIDE AND *VIRUCIDE AT 2 non-porous inanimate surfaces add 2 ounces of this

*VIRUCIDAL ACTIVITY - Lemon Disinfectant Neutral OUNCES PER GALLON. MODIFIED IN THE product per gallon of water, Apply solution with a cloth.

activity against HIV-1. HIV-2. Hepatitis B Virus (HBV). GALLON, MODIFIED IN THE PRESENCE OF 200 PPM solution for each use. Repeat application at weekly

wet all surfaces completely. Let air dry. Prepare a fresh

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

KEEP OUT OF REACH OF CHILDREN, CORROSIVE

Causes irreversible eve damage and skin burns. Do not

get in eyes, on skin or on clothing. May be fatal if absorbed

intervals or when mildew growth appears.

Cleaner - 64 when used on environmental, inanimate, PRESENCE OF 400 PPM HARD WATER AS CaCO3.

Hepatitis C Virus (HCV). Herpes Simplex Type 1 (causative HARD WATER AS CaCO3.

nfluenza A2/Hong Kong. Vaccinia. Rotavirus. Human called swine flu)

hard. non-porous surfaces exhibits effective virucidal THIS PRODUCT IS A FUNGICIDE AT 2 OUNCES PER

igent of fever blisters). Herpes Simplex Type 2 (genital). Kills Pandemic 2009 H1N1 Influenza A virus (formerly

Coronavirus (ATCC VR-740. Strain 229E). SARS Veterinary Practice/Animal Care/Anima

Associated Coronavirus, Rabies Virus, Paramyxovirus Laboratory/Zoos/Pet Shop/Kennels Disinfection

Mumps). Boying Viral Diarrhea Virus (BVDV), Directions; For cleaning and disinfecting the following

Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, hard non-porous surfaces; equipment not used for anima

lemon disinfectant CLEANER · DISINFECTANT · * VIRUCIDE · KILLS GERMS · DEODORIZEF **ACTIVE INGREDIENTS** Octyl decyl dimethyl ammonium chloride 1.6279 Dioctyl dimethyl ammonium chloride 0.814% Alkyl (50% C₁₄, 40% C₁₅, 10% C₁₆)

DANGER Have the product container or label with you when calling a poison control Lemon Disinfectant Neutral Cleaner - 64 deodorizes those heavily soiled areas, a pre-cleaning step is required, Apply NRS 384 Genotype USA300. Salmonella (enteritidis

center or doctor, or going for treatment. areas which generally are hard to keep fresh smelling, solution with a mop, cloth, sponge, hand pump trigger enterica. Ampicillin resistant Acinetobacter baumannii

KEEP OUT OF REACH OF CHILDREN

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20

immediately with plenty of water for 15-20 minutes. Call a poison control

center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air, If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, possible. Call a poison control center or doctor for further treatment advice You may contact 800/535-5053 for emergency medical treatment information

See side panels for additional precautionary statements. **NET CONTENTS: 1 GALLON (3.785 L)**

NOTE TO PHYSICIAN

EPA Reg. No. 1839-169-70925

EPA Est. No. 70925-CA-001

Chemcor Chemical Corporation 13770 Benson Ave., Chino, CA 91710

not breathe spray mist. For heavily soiled areas, a pre-cleaning step is required.

contamination from treated surfaces.

and their odors when used as directed.

manner inconsistent with its labeling

DIRECTIONS FOR USE

thoroughly. Allow to remain wet for 10 minutes, then Salmonella (choleraesuis) enterica, Staphylococcus Diluted product should be applied daily or more frequently remove excess liquid. For sprayer applications, spray 6-8 aureus, Staphylococcus aureus (clinical isolate), with heavy facility use. inches from surface, rub with brush, sponge or cloth. Do Bordetella bronchiseptica, Corynebacterium ammoniagenes, Enterobacter aerogenes, Enterobacter cloacae, Enterobacter cloacae (clinical isolate).

countertops, appliances, and tables, add 2 ounces of this Acinetobacter baumannii. Levofloxacin resistan cleaning step is required. Apply solution with a cloth. Acinetobacter baumannii. sponge or hand pump trigger sprayer so as to wet all FUNGICIDAL ACTIVITY - At the 2 ounce per gallor minutes, then remove excess liquid and rinse the surface fungicidal against the pathogenic fungi. Trichophytor

Enterococcus faecalis, Enterococcus faecalis (clinica

It is a violation of Federal Law to use this product in a surfaces thoroughly Allow the surface to remain wet for 10 dilution. Lemon Disinfectant Neutral Cleaner - 64 is Lemon Disinfectant Neutral Cleaner - 64 is not for use on with potable water. This product cannot be used to clean mentagrophytes (Athlete's Foot Fungus) and Candida

the following food contact surfaces: utensils, glassware albicans. Apply solution with a cloth, sponge or hand

critical and semi-critical medical device surfaces.

DISINFECTION - To disinfect inanimate, hard non- and dishes.

used with a mop and bucket, trigger sprayers or sponge.

This product is an economical concentrate that can be product per gallon of water. For heavily soiled areas, a pre- Acinetobacter baumannii and Bactrim resistan

neutral pH and will not dull high-gloss floor finishes with To disinfect restaurant food contact surfaces; Acinetobacter baumannii, Ciprofloxacin resistan

Lemon Disinfectant Neutral Cleaner - 64 is formulated to a area must be thoroughly rinsed with potable water.

processing plants, food service establishments, Directions; Remove gross filth prior to disinfection,

such as garbage storage areas, empty garbage bins and sprayer or low pressure coarse sprayer so as to wet all Cefazolin resistant Acinetobacter baumannii. Ceftazdime cans, toilet bowls, and other areas which are prone to surfaces thoroughly. Allow to remain wet for 10 minutes, resistant Acinetobacter baumannii, Ceftriaxone resistant then remove excess liquid. After use, all surfaces in the Acinetobacter baumannii. Gentamicin resistan

stainless steel surfaces, glazed porcelain, glazed ceramic and packaging materials must be removed from the area Methicillin resistant Staphylococcus aureus (CA-MRSA) tile, plastic surfaces, vanity tops, shower stalls, bathtubs, or carefully protected. For floors, walls, and storage areas, NRS 123 Genotype USA400. Community Associated

This product, when used as directed, is formulated to minutes and flush. disinfect the following hard non-porous inanimate To disinfect food processing premises: Before using (MRSA), Vancomycin intermediate resistan environmental surfaces; floors, walls, metal surfaces, Lemon Disinfectant Neutral Cleaner - 64, food products Staphylococcus aureus (VISA), Community Associated

Lemon Disinfectant Neutral Cleaner - 64 is a phosphate- BACTERICIDAL STABILITY OF USE-SOLUTION: isolate), Escherichia coli, Escherichia coli (clinica

free, pH neutral formulation designed to provide effective Prepare a fresh use-solution daily or more often if the use-jisolate). Escherichia coli O111:H8. Fusobacterium

cleaning, deodorizing, and disinfection for hospitals, solution becomes visibly dirty, contaminated or diluted, necrophorum, Klebsiella pneumoniae subsp

nursing homes, households, homes, schools, food Non-Acid Toilet Bowl Disinfection/Cleaner pneumoniae, Lactobacillus casei subsp. rhamnosus

restaurants, transportation terminals, office buildings, From concentrate; Add 2 ounces per gallon to the toilet vulgaris, Proteus mirabilis ATCC 9921, Proteus mirabilis

manufacturing facilities, lodging establishments, hotels, bowl and mix. Brush thoroughly over exposed surfaces ATCC 25933, Salmonella (paratyphi B) enterica

retail business, and athletic/recreation facilities, sports and under the rim, allow to stand for 10 minutes and flush. Salmonella (typhi) enterica, Salmonella (

stadiums, amphitheaters and convention centers where From use-solution; Empty toilet bowl or urinal and apply enterica. Salmonella (pullorum) enterica. Serratia

cloth, mop, sponge or sprayer, allow to stand for 10 isolate). Vancomycin resistant Enterococcus faecalis (VRF). Methicillin resistant Stanhylococcus aureus

neutral disinfectant that is effective against a broad For heavy duty use: Empty toilet bowl or urinal and epidermidis (clinical isolate), Streptococcus pyogenes spectrum of bacteria, is virucidal* (including HIV-1, HIV-2, apply 8 ounces per gallon use-solution to exposed (Clinical - Flesh Eating Strain BIRD M3), Streptococcus HBV and HCV) and inhibits the growth of mold and mildew surfaces, brush or swab thoroughly under the rim with a pyogenes Group A. Xanthamonas maltophilia (clinica

add 2 ounces of this product per gallon of water. For Methicillin resistant Staphylococcus aureus (CA-MRSA)

housekeeping is of prime importance in controlling cross- 2 ounces per gallon use-solution to exposed surfaces, marcescens, Shigella sonnei, Shigella flexneri Type 2b brush or swab thoroughly under the rim with a cloth, mop. Shigella dysenteriae. Staphylococcus aureus subsp Lemon Disinfectant Neutral Cleaner - 64 is a one step sponge or sprayer, allow to stand for 10 minutes and flush, aureus, Staphylococcus epidermidis. Staphylococcus

Listeria monocytogenes, Pasteurella multocida, Proteus

Acinetobacter baumannii. Tobramycin resistan

pump trigger sprayer to hard, non-porous surfaces found porous surfaces add 2 ounces of Lemon Disinfectant BACTERICIDAL ACTIVITY - At the 2 ounce per gallon in bathrooms, shower stalls, locker rooms, exercise Neutral Cleaner - 64 per gallon of water. Apply solution dilution Lemon Disinfectant Neutral Cleaner - 64 facilities or other clean, hard non-porous surfaces with a mop, cloth, sponge, hand pump trigger sprayer or demonstrates effective disinfectant activity against the commonly contacted by bare feet. Allow the surface to low pressure coarse sprayer so as to wet all surfaces organisms: Pseudomonas aeruginosa PRD-10, remain wet for 10 minutes, then remove excess liquid